

Parabolic Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Ivan Ip

Hong Kong University of Science and Technology

March 16, 2021

Geometry, Number Theory, and Representation Theory Seminar,
University of Alberta

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Definition of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$

Definition

$\mathcal{U}_q(\mathfrak{sl}_2) =$ Hopf-algebra $\langle E, F, K^{\pm 1} \rangle$ over $\mathbb{C}(q)$ such that

$$KE = q^2 EK, \quad KF = q^{-2} FK, \quad [E, F] = \frac{K - K^{-1}}{q - q^{-1}}$$

Coproduct:

$$\begin{aligned} \Delta(E) &= 1 \otimes E + E \otimes K, & \Delta(F) &= F \otimes 1 + K^{-1} \otimes F \\ \Delta(K) &= K \otimes K \end{aligned}$$

(Also counit ϵ , antipode S)

$$\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R})): \quad (|q| = 1)$$

$$E^* = E, \quad F^* = F, \quad K^* = K$$

Definition of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$

Definition

$\mathcal{U}_q(\mathfrak{sl}_2) =$ Hopf-algebra $\langle E, F, K^{\pm 1} \rangle$ over $\mathbb{C}(q)$ such that

$$KE = q^2 EK, \quad KF = q^{-2} FK, \quad [E, F] = \frac{K - K^{-1}}{q - q^{-1}}$$

Coproduct:

$$\begin{aligned} \Delta(E) &= 1 \otimes E + E \otimes K, & \Delta(F) &= F \otimes 1 + K^{-1} \otimes F \\ \Delta(K) &= K \otimes K \end{aligned}$$

(Also counit ϵ , antipode S)

$$\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R})): \quad (|q| = 1)$$

$$E^* = E, \quad F^* = F, \quad K^* = K$$

Definition of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Definition

$\mathcal{U}_q(\mathfrak{g}) = \text{Hopf-algebra } \langle E_i, F_i, K_i^{\pm 1} \rangle_{i \in I} \text{ over } \mathbb{C}(q) \text{ such that}$

Cartan matrix

$$K_i E_j = q^{a_{ij}} E_j K_i, \quad K_i F_j = q^{-a_{ij}} F_j K_i, \quad [E_i, F_j] = \delta_{ij} \frac{K_i - K_i^{-1}}{q - q^{-1}}$$

+ Serre relations.

Coproduct:

$$\Delta(E_i) = 1 \otimes E_i + E_i \otimes K_i, \quad \Delta(F_i) = F_i \otimes 1 + K_i^{-1} \otimes F_i$$

$$\Delta(K_i) = K_i \otimes K_i$$

(Also counit ϵ , antipode S)

$\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}}): \quad (|q| = 1) \quad q^* := \bar{q}$

$$E_i^* = E_i, \quad F_i^* = F_i, \quad K_i^* = K_i$$

Definition of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Definition

$\mathcal{D}_q(\mathfrak{g}) =$ Drinfeld's Double: $\langle E_i, F_i, K_i^{\pm 1}, K_i'^{\pm 1} \rangle_{i \in I}$
of $\mathcal{U}_q(\mathfrak{g})$

$$K_i E_j = q^{a_{ij}} E_j K_i, \quad K_i F_j = q^{-a_{ij}} F_j K_i, \quad [E_i, F_j] = \delta_{ij} \frac{K_i - K_i'}{q - q^{-1}}$$

+ Serre relations + Similar for K_i'

Coproduct:

$$\begin{aligned} \Delta(E_i) &= 1 \otimes E_i + E_i \otimes K_i, & \Delta(F_i) &= F_i \otimes 1 + K_i' \otimes F_i \\ \Delta(K_i) &= K_i \otimes K_i, & \Delta(K_i') &= K_i \otimes K_i' \end{aligned}$$

(Also counit ϵ , antipode S)

$$\mathcal{U}_q(\mathfrak{g}) = \mathcal{D}_q(\mathfrak{g}) / \langle K_i K_i' = 1 \rangle_{i \in I}$$

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations by positive operators on Hilbert space.
- Generalization of Teschner's representations of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$
 - Closure under taking tensor product A_n : [Schrader-Shapiro 2018]
 - Braiding structure [I. 2012]
 - Peter-Weyl Theorem A_n : [I.-Schrader-Shapiro 2020]
- = “Quantization of principal series representations”
- Constructed for all semisimple Lie types.

Construction:

- Lusztig's total positive space $L^2((G/B)_{>0}) \simeq L^2(\mathbb{R}_{>0}^{N=\ell(w_0)})$
- Mellin transformation: $L^2(\mathbb{R}_{>0}^N) \simeq L^2(\mathbb{R}^N)$
- $\mathcal{U}(\mathfrak{g})$ differential operator \leadsto finite difference operator
- Quantization \leadsto positive operators $\mathbf{e}_i, \mathbf{f}_i, K_i \in \mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations by positive operators on Hilbert space.
- Generalization of Teschner's representations of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$
 - Closure under taking tensor product A_n : [Schrader-Shapiro 2018]
 - Braiding structure [I. 2012]
 - Peter-Weyl Theorem A_n : [I.-Schrader-Shapiro 2020]
- = “Quantization of principal series representations”
- Constructed for all semisimple Lie types.

Construction:

- Lusztig's total positive space $L^2((G/B)_{>0}) \simeq L^2(\mathbb{R}_{>0}^{N=\ell(w_0)})$
- Mellin transformation: $L^2(\mathbb{R}_{>0}^N) \simeq L^2(\mathbb{R}^N)$
- $\mathcal{U}(\mathfrak{g})$ differential operator \leadsto finite difference operator
- Quantization \leadsto positive operators $\mathbf{e}_i, \mathbf{f}_i, K_i \in \mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations by positive operators on Hilbert space.
- Generalization of Teschner's representations of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$
 - Closure under taking tensor product A_n : [Schrader-Shapiro 2018]
 - Braiding structure [I. 2012]
 - Peter-Weyl Theorem A_n : [I.-Schrader-Shapiro 2020]
- = “Quantization of principal series representations”
- Constructed for all semisimple Lie types.

Construction:

- Lusztig's total positive space $L^2((G/B)_{>0}) \simeq L^2(\mathbb{R}_{>0}^{N=\ell(w_0)})$
- Mellin transformation: $L^2(\mathbb{R}_{>0}^N) \simeq L^2(\mathbb{R}^N)$
- $\mathcal{U}(\mathfrak{g})$ differential operator \leadsto finite difference operator
- Quantization \leadsto positive operators $e_i, f_i, K_i \in \mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations by positive operators on Hilbert space.
- Generalization of Teschner's representations of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$
- Closure under taking tensor product A_n : [Schrader-Shapiro 2018]
- Braiding structure [I. 2012]
- Peter-Weyl Theorem A_n : [I.-Schrader-Shapiro 2020]
- = "Quantization of principal series representations"
- Constructed for all semisimple Lie types.

Construction:

- Lusztig's total positive space $L^2((G/B)_{>0}) \simeq L^2(\mathbb{R}_{>0}^{N=\ell(w_0)})$
- Mellin transformation: $L^2(\mathbb{R}_{>0}^N) \simeq L^2(\mathbb{R}^N)$
- $\mathcal{U}(\mathfrak{g})$ differential operator \leadsto finite difference operator
- Quantization \leadsto positive operators $e_i, f_i, K_i \in \mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations by positive operators on Hilbert space.
- Generalization of Teschner's representations of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$
 - Closure under taking tensor product A_n : [Schrader-Shapiro 2018]
 - Braiding structure [I. 2012]
 - Peter-Weyl Theorem A_n : [I.-Schrader-Shapiro 2020]
- = “Quantization of principal series representations”
- Constructed for all semisimple Lie types.

Construction:

- Lusztig's total positive space $L^2((G/B)_{>0}) \simeq L^2(\mathbb{R}_{>0}^{N=\ell(w_0)})$
- Mellin transformation: $L^2(\mathbb{R}_{>0}^N) \simeq L^2(\mathbb{R}^N)$
- $\mathcal{U}(\mathfrak{g})$ differential operator \leadsto finite difference operator
- Quantization \leadsto positive operators $e_i, f_i, K_i \in \mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations by positive operators on Hilbert space.
- Generalization of Teschner's representations of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$
 - Closure under taking tensor product A_n : [Schrader-Shapiro 2018]
 - Braiding structure [I. 2012]
 - Peter-Weyl Theorem A_n : [I.-Schrader-Shapiro 2020]
- = “Quantization of principal series representations”
- Constructed for all semisimple Lie types.

Construction:

- Lusztig's total positive space $L^2((G/B)_{>0}) \simeq L^2(\mathbb{R}_{>0}^{N=\ell(w_0)})$
- Mellin transformation: $L^2(\mathbb{R}_{>0}^N) \simeq L^2(\mathbb{R}^N)$
- $\mathcal{U}(\mathfrak{g})$ differential operator \leadsto finite difference operator
- Quantization \leadsto positive operators $e_i, f_i, K_i \in \mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations by positive operators on Hilbert space.
- Generalization of Teschner's representations of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$
 - Closure under taking tensor product A_n : [Schrader-Shapiro 2018]
 - Braiding structure [I. 2012]
 - Peter-Weyl Theorem A_n : [I.-Schrader-Shapiro 2020]
- = “Quantization of principal series representations”
- Constructed for all semisimple Lie types.

Construction:

- Lusztig's total positive space $L^2((G/B)_{>0}) \simeq L^2(\mathbb{R}_{>0}^{N=\ell(w_0)})$
- Mellin transformation: $L^2(\mathbb{R}_{>0}^N) \simeq L^2(\mathbb{R}^N)$
- $\mathcal{U}(\mathfrak{g})$ differential operator \leadsto finite difference operator
- Quantization \leadsto positive operators $e_i, f_i, K_i \in \mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations by positive operators on Hilbert space.
- Generalization of Teschner's representations of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$
 - Closure under taking tensor product A_n : [Schrader-Shapiro 2018]
 - Braiding structure [I. 2012]
 - Peter-Weyl Theorem A_n : [I.-Schrader-Shapiro 2020]
- = “Quantization of **principal series representations**”
- Constructed for all semisimple Lie types.

Construction:

- Lusztig's **total positive** space $L^2((G/B)_{>0}) \simeq L^2(\mathbb{R}_{>0}^{N=\ell(w_0)})$
- Mellin transformation: $L^2(\mathbb{R}_{>0}^N) \simeq L^2(\mathbb{R}^N)$
- $\mathcal{U}(\mathfrak{g})$ differential operator \rightsquigarrow finite difference operator
- Quantization \rightsquigarrow **positive operators** $e_i, f_i, K_i \in \mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations by positive operators on Hilbert space.
- Generalization of Teschner's representations of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$
 - Closure under taking tensor product A_n : [Schrader-Shapiro 2018]
 - Braiding structure [I. 2012]
 - Peter-Weyl Theorem A_n : [I.-Schrader-Shapiro 2020]
- = “Quantization of principal series representations”
- Constructed for all semisimple Lie types.

Construction:

- Lusztig's total positive space $L^2((G/B)_{>0}) \simeq L^2(\mathbb{R}_{>0}^{N=\ell(w_0)})$
- Mellin transformation: $L^2(\mathbb{R}_{>0}^N) \simeq L^2(\mathbb{R}^N)$
- $\mathcal{U}(\mathfrak{g})$ differential operator \rightsquigarrow finite difference operator
- Quantization \rightsquigarrow positive operators $e_i, f_i, K_i \in \mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations by positive operators on Hilbert space.
- Generalization of Teschner's representations of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$
 - Closure under taking tensor product A_n : [Schrader-Shapiro 2018]
 - Braiding structure [I. 2012]
 - Peter-Weyl Theorem A_n : [I.-Schrader-Shapiro 2020]
- = “Quantization of principal series representations”
- Constructed for all semisimple Lie types.

Construction:

- Lusztig's total positive space $L^2((G/B)_{>0}) \simeq L^2(\mathbb{R}_{>0}^{N=\ell(w_0)})$
- Mellin transformation: $L^2(\mathbb{R}_{>0}^N) \simeq L^2(\mathbb{R}^N)$
- $\mathcal{U}(\mathfrak{g})$ differential operator \leadsto finite difference operator
- Quantization \leadsto positive operators $e_i, f_i, K_i \in \mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Research program started in [Frenkel-I. (2012)]

- Representations by positive operators on Hilbert space.
- Generalization of Teschner's representations of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$
 - Closure under taking tensor product A_n : [Schrader-Shapiro 2018]
 - Braiding structure [I. 2012]
 - Peter-Weyl Theorem A_n : [I.-Schrader-Shapiro 2020]
- = "Quantization of principal series representations"
- Constructed for all semisimple Lie types.

Construction:

- Lusztig's total positive space $L^2((G/B)_{>0}) \simeq L^2(\mathbb{R}_{>0}^{N=\ell(w_0)})$
- Mellin transformation: $L^2(\mathbb{R}_{>0}^N) \simeq L^2(\mathbb{R}^N)$
- $\mathcal{U}(\mathfrak{g})$ differential operator \leadsto finite difference operator
- Quantization \leadsto positive operators $\mathbf{e}_i, \mathbf{f}_i$, $K_i \in \mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

$$e_i = \left(\frac{i}{i-i'}\right)' E_i$$

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Rescale generators by $(q = e^{\pi i b^2}, b \in (0, 1)) \searrow \mathbb{Q}$

$$\mathbf{e}_k = -i(q - q^{-1})E_k, \quad \mathbf{f}_k = -i(q - q^{-1})F_k$$

Theorem (I. (2012))

There exists a family of irreducible representations \mathcal{P}_λ of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$:

- Parametrized by $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$
- *Positivity*: $\{\mathbf{e}_i, \mathbf{f}_i, K_i\}$ are represented by positive, essentially self-adjoint (unbounded) operators on $L^2(\mathbb{R}^N)$
- $\mathbf{e}_i, \mathbf{f}_i, K_i$ are expressed in terms of Laurent polynomials of $\{e^{\pi b x_k}, e^{2\pi b p_k}\}_{k=1}^N$
- Characterized by *modular double* structure (*Langland's duality*)
- One can recover *any* finite dimensional irreducible representations of $\mathcal{U}_q(\mathfrak{g})$ by appropriate *analytic continuation* on λ .

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Rescale generators by $(q = e^{\pi i b^2}, b \in (0, 1))$

$$\mathbf{e}_k = -i(q - q^{-1})E_k, \quad \mathbf{f}_k = -i(q - q^{-1})F_k$$

Theorem (I. (2012))

There exists a family of irreducible representations \mathcal{P}_λ of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$:

- Parametrized by $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$
- *Positivity:* $\{\mathbf{e}_i, \mathbf{f}_i, K_i\}$ are represented by positive, essentially self-adjoint (unbounded) operators on $L^2(\mathbb{R}^N)$
- $\mathbf{e}_i, \mathbf{f}_i, K_i$ are expressed in terms of Laurent polynomials of $\{e^{\pi b x_k}, e^{2\pi b p_k}\}_{k=1}^N$
- Characterized by *modular double* structure (*Langland's duality*)
- One can recover *any* finite dimensional irreducible representations of $\mathcal{U}_q(\mathfrak{g})$ by appropriate *analytic continuation* on λ .

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Rescale generators by $(q = e^{\pi i b^2}, b \in (0, 1))$

$$\mathbf{e}_k = -i(q - q^{-1})E_k, \quad \mathbf{f}_k = -i(q - q^{-1})F_k$$

Theorem (I. (2012))

There exists a family of irreducible representations \mathcal{P}_λ of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$:

- Parametrized by $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$
- *Positivity*: $\{\mathbf{e}_i, \mathbf{f}_i, K_i\}$ are represented by positive, essentially self-adjoint (unbounded) operators on $L^2(\mathbb{R}^N)$
- $\mathbf{e}_i, \mathbf{f}_i, K_i$ are expressed in terms of Laurent polynomials of $\{e^{\pi b x_k}, e^{2\pi b p_k}\}_{k=1}^N$
- Characterized by *modular double* structure (*Langland's duality*)
- One can recover *any* finite dimensional irreducible representations of $\mathcal{U}_q(\mathfrak{g})$ by appropriate *analytic continuation* on λ .

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Rescale generators by $(q = e^{\pi i b^2}, b \in (0, 1))$

$$\mathbf{e}_k = -i(q - q^{-1})E_k, \quad \mathbf{f}_k = -i(q - q^{-1})F_k$$

Theorem (I. (2012))

There exists a family of irreducible representations \mathcal{P}_λ of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$:

- Parametrized by $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$
- **Positivity:** $\{\mathbf{e}_i, \mathbf{f}_i, K_i\}$ are represented by positive, essentially self-adjoint (unbounded) operators on $L^2(\mathbb{R}^N)$
- $\mathbf{e}_i, \mathbf{f}_i, K_i$ are expressed in terms of Laurent polynomials of $\{e^{\pi b x_k}, e^{2\pi b p_k}\}_{k=1}^N$
- Characterized by **modular double** structure (**Langland's duality**)
- One can recover **any** finite dimensional irreducible representations of $\mathcal{U}_q(\mathfrak{g})$ by appropriate **analytic continuation** on λ .

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Rescale generators by $(q = e^{\pi i b^2}, b \in (0, 1))$

$$\mathbf{e}_k = -i(q - q^{-1})E_k, \quad \mathbf{f}_k = -i(q - q^{-1})F_k$$

Theorem (I. (2012))

There exists a family of irreducible representations \mathcal{P}_λ of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$:

- Parametrized by $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$
- **Positivity:** $\{\mathbf{e}_i, \mathbf{f}_i, K_i\}$ are represented by positive, essentially self-adjoint (unbounded) operators on $L^2(\mathbb{R}^N)$
- $\mathbf{e}_i, \mathbf{f}_i, K_i$ are expressed in terms of Laurent polynomials of $\{e^{\pi b x_k}, e^{2\pi b p_k}\}_{k=1}^N$

$p = \frac{1}{2\pi i} \frac{\partial}{\partial x}$

$e^{2\pi b p} f(x) = f(x - ib)$
- Characterized by *modular double* structure (Langland's duality)
- One can recover **any** finite dimensional irreducible representations of $\mathcal{U}_q(\mathfrak{g})$ by appropriate *analytic continuation* on λ .

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Rescale generators by $(q = e^{\pi i b^2}, b \in (0, 1))$

$$\mathbf{e}_k = -i(q - q^{-1})E_k, \quad \mathbf{f}_k = -i(q - q^{-1})F_k$$

Theorem (I. (2012))

There exists a family of irreducible representations \mathcal{P}_λ of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$:

- Parametrized by $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$
- **Positivity:** $\{\mathbf{e}_i, \mathbf{f}_i, K_i\}$ are represented by positive, essentially self-adjoint (unbounded) operators on $L^2(\mathbb{R}^N)$
- $\mathbf{e}_i, \mathbf{f}_i, K_i$ are expressed in terms of Laurent polynomials of $\{e^{\pi b x_k}, e^{2\pi b p_k}\}_{k=1}^N$
- Characterized by **modular double** structure (**Langland's duality**)
- One can recover **any** finite dimensional irreducible representations of $\mathcal{U}_q(\mathfrak{g})$ by appropriate **analytic continuation** on λ .

$b \leftrightarrow b^{-1}$

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Rescale generators by $(q = e^{\pi i b^2}, b \in (0, 1))$

$$\mathbf{e}_k = -i(q - q^{-1})E_k, \quad \mathbf{f}_k = -i(q - q^{-1})F_k$$

Theorem (I. (2012))

There exists a family of irreducible representations \mathcal{P}_λ of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$:

- Parametrized by $\lambda \in \mathbb{R}_{\geq 0} P^+ \simeq \mathbb{R}_{\geq 0}^{n=\text{rank } \mathfrak{g}}$
- *Positivity*: $\{\mathbf{e}_i, \mathbf{f}_i, K_i\}$ are represented by positive, essentially self-adjoint (unbounded) operators on $L^2(\mathbb{R}^N)$
- $\mathbf{e}_i, \mathbf{f}_i, K_i$ are expressed in terms of Laurent polynomials of $\{e^{\pi b x_k}, e^{2\pi b p_k}\}_{k=1}^N$
- Characterized by *modular double* structure (*Langland's duality*)
- One can recover *any* finite dimensional irreducible representations of $\mathcal{U}_q(\mathfrak{g})$ by appropriate *analytic continuation* on λ .

Example: $\mathcal{U}_q(\mathfrak{sl}_3)$

Coordinates on $(G/B)_{>0}$:

$x_1(a)$ $x_2(b)$ $x_1(c)$

$$\begin{pmatrix} 1 & a & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & c & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & t & 1 \end{pmatrix} \quad a, b, c > 0$$

e^{tF_2}

$$= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & \frac{t}{1+bt} & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1+bt & 0 \\ 0 & 0 & (1+bt)^{-1} \end{pmatrix} \begin{pmatrix} 1 & a+abt & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & \frac{b}{1+bt} \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & c & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$e^{tF_2} \cdot f(a, b, c) = (1+bt)^{2\lambda} f(a+abt, \frac{b}{1+bt}, c), \quad \lambda \in \mathbb{R}_{\geq 0}$$

$$F_2 := \left. \frac{d}{dt} e^{tF_2} \right|_{t=0} = ab \frac{\partial}{\partial a} - b^2 \frac{\partial}{\partial b} + b\lambda$$

Example: $\mathcal{U}_q(\mathfrak{sl}_3)$ Coordinates on $(G/B)_{>0}$:

$$\begin{pmatrix} 1 & a & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & c & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & t & 1 \end{pmatrix} \quad a, b, c > 0$$

$$= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & t & 1+bt \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1+bt & 0 \\ 0 & 0 & (1+bt)^{-1} \end{pmatrix} \begin{pmatrix} 1 & a+abt & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & \frac{b}{1+bt} \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & c & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

u^- T x_λ x_1 x_2 x_1

$$e^{tF_2} \cdot f(a, b, c) = (1+bt)^{2\lambda} f(a+abt, \frac{b}{1+bt}, c), \quad \lambda \in \mathbb{R}_{\geq 0}$$

$$F_2 := \left. \frac{d}{dt} e^{tF_2} \right|_{t=0} = ab \frac{\partial}{\partial a} - b^2 \frac{\partial}{\partial b} + b\lambda$$

Example: $\mathcal{U}_q(\mathfrak{sl}_3)$

Coordinates on $(G/B)_{>0}$:

$$\begin{pmatrix} 1 & a & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & c & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & t & 1 \end{pmatrix} \quad a, b, c > 0$$

$$= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & \frac{t}{1+bt} & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1+bt & 0 \\ 0 & 0 & (1+bt)^{-1} \end{pmatrix} \begin{pmatrix} 1 & a+abt & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & \frac{b}{1+bt} \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & c & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$e^{tF_2} \cdot f(a, b, c) = (1+bt)^{2\lambda} f(a+abt, \frac{b}{1+bt}, c), \quad \lambda \in \mathbb{R}_{\geq 0}$$

$$F_2 := \left. \frac{d}{dt} e^{tF_2} \right|_{t=0} = ab \frac{\partial}{\partial a} - b^2 \frac{\partial}{\partial b} + b\lambda$$

Example: $\mathcal{U}_q(\mathfrak{sl}_3)$

Coordinates on $(G/B)_{>0}$:

$$\begin{pmatrix} 1 & a & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & c & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & t & 1 \end{pmatrix} \quad a, b, c > 0$$

$$= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & \frac{t}{1+bt} & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1+bt & 0 \\ 0 & 0 & (1+bt)^{-1} \end{pmatrix} \begin{pmatrix} 1 & a+abt & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & \frac{b}{1+bt} \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & c & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$e^{tF_2} \cdot f(a, b, c) = (1+bt)^{2\lambda} f(a+abt, \frac{b}{1+bt}, c), \quad \lambda \in \mathbb{R}_{\geq 0}$$

$$F_2 := \left. \frac{d}{dt} e^{tF_2} \right|_{t=0} = ab \frac{\partial}{\partial a} - b^2 \frac{\partial}{\partial b} + b\lambda$$

Example: $\mathcal{U}_q(\mathfrak{sl}_3)$

$$F_2 = ab \frac{\partial}{\partial a} - b^2 \frac{\partial}{\partial b} + b\lambda$$

(Formal) Mellin transform: $\mathcal{F}(u, v, w) := \int f(a, b, c) a^u b^v c^w da db dc$

$$F_2 : \mathcal{F}(u, v, w) \mapsto (2\lambda + u - v + 1) \mathcal{F}(u, v - 1, w)$$

Quantum Twist ($n \mapsto [n]_q + \text{“Wick’s rotation”}$)

$$F_2 := \left(\frac{i}{q - q^{-1}} \right) \left(e^{\pi b(2\lambda + u - v + 2p_v)} + e^{\pi b(-2\lambda - u + v + 2p_v)} \right)$$

Example: $\mathcal{U}_q(\mathfrak{sl}_3)$

$$F_2 = ab \frac{\partial}{\partial a} - b^2 \frac{\partial}{\partial b} + b\lambda$$

(Formal) Mellin transform: $\mathcal{F}(u, v, w) := \int f(a, b, c) a^u b^v c^w da db dc$

$$F_2 : \mathcal{F}(u, v, w) \mapsto (2\lambda + u - v + 1) \mathcal{F}(u, v - 1, w)$$

Quantum Twist ($n \mapsto [n]_q$ + “Wick’s rotation”)

$$F_2 := \left(\frac{i}{q - q^{-1}} \right) \left(e^{\pi b(2\lambda + u - v + 2p_v)} + e^{\pi b(-2\lambda - u + v + 2p_v)} \right)$$

Example: $\mathcal{U}_q(\mathfrak{sl}_3)$

$$F_2 = ab \frac{\partial}{\partial a} - b^2 \frac{\partial}{\partial b} + b\lambda$$

(Formal) Mellin transform: $\mathcal{F}(u, v, w) := \int f(a, b, c) a^u b^v c^w da db dc$

$$F_2 : \mathcal{F}(u, v, w) \mapsto \left[(2\lambda + u - v + 1) \right]_q \mathcal{F}(u, v - 1, w)$$

$[n] = \frac{q^n - q^{-n}}{q - q^{-1}}$

Quantum Twist ($n \mapsto [n]_q$ + “Wick’s rotation”) $u \mapsto -\frac{i}{b} u$

$$F_2 := \left(\frac{i}{q - q^{-1}} \right) \left(e^{\pi b(2\lambda + u - v + 2p_v)} + e^{\pi b(-2\lambda - u + v + 2p_v)} \right)$$

The goal of this talk

Definition

Parabolic positive representations is a new family of positive representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ based on quantizing the parabolic induction representations on $L^2((G/P)_{>0})$, where $P \subset G$ is a parabolic subgroup.

- It answers some combinatorial mysteries of quantum group embedding (cluster realization)
- Gives a new realization of the evaluation module of $\mathcal{U}_q(\widehat{\mathfrak{sl}}_n)$.

The goal of this talk

Definition

Parabolic positive representations is a new family of positive representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ based on quantizing the parabolic induction representations on $L^2((G/P)_{>0})$, where $P \subset G$ is a parabolic subgroup.

- It answers some **combinatorial mysteries** of quantum group embedding (cluster realization)
- Gives a new realization of the **evaluation module** of $\mathcal{U}_q(\widehat{\mathfrak{sl}}_n)$.

Quantum Cluster Variety

Quantum Torus Algebra

“Quantization of cluster \mathcal{X} variety” [Fock-Goncharov]

Definition

Seed $\mathbf{Q} = (Q, Q_0, B)$:

- $Q = \text{nodes (finite set)}$
- $Q_0 \subset Q = \text{frozen nodes}$
- $B = (b_{ij})$ exchange matrix ($|Q| \times |Q|$, skew-symmetric, $\frac{1}{2}\mathbb{Z}$ -valued)

Quantum torus algebra $\mathcal{X}_q^{\mathbf{Q}} = \text{algebra generated by } \{X_i\}_{i \in Q} \text{ over } \mathbb{C}[q]$
such that

$$X_i X_j = q^{-2b_{ij}} X_j X_i$$

$X_i = \text{quantum cluster variables}$

Exchange Matrix $B \rightsquigarrow$ Quiver.



$$X_i X_j = q^{-2} X_j X_i,$$

Quantum Torus Algebra

“Quantization of cluster \mathcal{X} variety” [Fock-Goncharov]

Definition

Seed $\mathbf{Q} = (Q, Q_0, B)$:

- $Q = \text{nodes (finite set)}$
- $Q_0 \subset Q = \text{frozen nodes}$
- $B = (b_{ij})$ exchange matrix ($|Q| \times |Q|$, skew-symmetric, $\frac{1}{2}\mathbb{Z}$ -valued)

Quantum torus algebra $\mathcal{X}_q^{\mathbf{Q}} = \text{algebra generated by } \{X_i\}_{i \in Q} \text{ over } \mathbb{C}[q]$
such that

$$X_i X_j = q^{-2b_{ij}} X_j X_i$$

$X_i = \text{quantum cluster variables}$

Exchange Matrix $B \rightsquigarrow$ Quiver.



$$X_i X_j = q^{-2} X_j X_i,$$

Quantum Torus Algebra

“Quantization of cluster \mathcal{X} variety” [Fock-Goncharov]

Definition

Seed $\mathbf{Q} = (Q, Q_0, B)$:

- $Q = \text{nodes (finite set)}$
- $Q_0 \subset Q = \text{frozen nodes}$
- $B = (b_{ij})$ exchange matrix ($|Q| \times |Q|$, skew-symmetric, $\frac{1}{2}\mathbb{Z}$ -valued)

Quantum torus algebra $\mathcal{X}_q^{\mathbf{Q}} = \text{algebra generated by } \{X_i\}_{i \in Q} \text{ over } \mathbb{C}[q]$
such that

$$X_i X_j = q^{-2b_{ij}} X_j X_i$$

$X_i = \text{quantum cluster variables}$

Exchange Matrix $B \rightsquigarrow$ Quiver.



$$X_i X_j = q^{-2} X_j X_i,$$

Quantum Torus Algebra

“Quantization of cluster \mathcal{X} variety” [Fock-Goncharov]

Definition

Seed $\mathbf{Q} = (Q, Q_0, B)$:


- $Q = \text{nodes (finite set)}$
- $Q_0 \subset Q = \text{frozen nodes}$
- $B = (b_{ij})$ exchange matrix ($|Q| \times |Q|$, skew-symmetric, $\frac{1}{2}\mathbb{Z}$ -valued)

Quantum torus algebra $\mathcal{X}_q^{\mathbf{Q}} = \text{algebra generated by } \{X_i\}_{i \in Q} \text{ over } \mathbb{C}[q]$
such that

$$X_i X_j = q^{-2b_{ij}} X_j X_i$$

$X_i = \text{quantum cluster variables}$

Exchange Matrix $B \rightsquigarrow$ Quiver.



$$\boxed{i} \cdots \cdots \rightarrow \boxed{j} \quad X_i X_j = q^{-1} X_j X_i$$

$$X_i X_j = q^{-2} X_j X_i,$$

Quantum Torus Algebra

“Quantization of cluster \mathcal{X} variety” [Fock-Goncharov]

Definition

Seed $\mathbf{Q} = (Q, Q_0, B)$:

- $\Lambda_{\mathbf{Q}} = \mathbb{Z}$ -Lattice with basis $\{e_i\}_{i \in Q}$
- $(-, -)$ skew-symmetric form, $(e_i, e_j) := b_{ij}$.


Quantum torus algebra $\mathcal{X}_q^{\mathbf{Q}}$ = algebra generated by $\{X_{\lambda}\}_{\lambda \in \Lambda_{\mathbf{Q}}}$ over $\mathbb{C}[q^{\frac{1}{2}}]$ such that

$$X_{\lambda+\mu} = q^{(\lambda, \mu)} X_{\lambda} X_{\mu}$$

$$X_i := X_{e_i}, \quad X_{i_1, i_2, \dots, i_k} := X_{e_{i_1} + e_{i_2} + \dots + e_{i_k}}$$

Exchange Matrix $B \rightsquigarrow$ Quiver.

$$q^{X_i X_j} =: X_{i,j}.$$



$$X_i X_j = q^{-2} X_j X_i,$$

Quantum Cluster Mutations

$\mathbf{T}_q^{\mathbf{Q}}$:= (non-commutative) field of fractions of $\mathcal{X}_q^{\mathbf{Q}}$.

Cluster mutation μ_k induces $\mu_k^q : \mathbf{T}_q^{\mathbf{Q}'} \longrightarrow \mathbf{T}_q^{\mathbf{Q}}$:

$$\mu_k^q(\widehat{X}_i) := \begin{cases} X_k^{-1} & i = k \\ X_i \prod_{r=1}^{|b_{ki}|} (1 + q_i^{2r-1} X_k) & i \neq k, b_{ki} < 0 \\ X_i \prod_{r=1}^{b_{ki}} (1 + q_i^{2r-1} X_k^{-1})^{-1} & i \neq k, b_{ki} > 0 \end{cases}$$

Can be rewritten as

$$\mu_k^q = \mu_k^{\#} \circ \mu_k'$$

$$\mu_k'(\widehat{X}_i) := \begin{cases} X_k^{-1} & i = k \\ X_i & i \neq k, b_{ki} < 0 \\ q_i^{b_{ik}b_{ki}} X_i X_k^{b_{ik}} & i \neq k, b_{ki} > 0 \end{cases}$$

$$\mu_k^{\#} := Ad_{\Psi_q(X_k)}$$

Ψ_q = quantum dilogarithm

Quantum Cluster Mutations

$\mathbf{T}_q^{\mathbf{Q}}$:= (non-commutative) field of fractions of $\mathcal{X}_q^{\mathbf{Q}}$.

Cluster mutation μ_k induces $\mu_k^q : \mathbf{T}_q^{\mathbf{Q}'} \longrightarrow \mathbf{T}_q^{\mathbf{Q}}$:

$$\mu_k^q(\widehat{X}_i) := \begin{cases} X_k^{-1} & i = k \\ X_i \prod_{r=1}^{|b_{ki}|} (1 + q_i^{2r-1} X_k) & i \neq k, b_{ki} < 0 \\ X_i \prod_{r=1}^{b_{ki}} (1 + q_i^{2r-1} X_k^{-1})^{-1} & i \neq k, b_{ki} > 0 \end{cases}$$

Can be rewritten as

$$\mu_k^q = \mu_k^{\#} \circ \mu'_k$$

$$\mu'_k(\widehat{X}_i) := \begin{cases} X_k^{-1} & i = k \\ X_i & i \neq k, b_{ki} < 0 \\ q_i^{b_{ik}b_{ki}} X_i X_k^{b_{ik}} & i \neq k, b_{ki} > 0 \end{cases}$$

$$\mu_k^{\#} := Ad_{\Psi_q(X_k)}$$

Ψ_q = quantum dilogarithm (non compact version \Rightarrow unitary transform)

Polarization of $\mathcal{X}_q^{\mathbf{Q}}$

Recall $q = e^{\pi i b^2}$ such that $|q| = 1$.

Definition

A *polarization* of $\mathcal{X}_q^{\mathbf{Q}}$ is a choice of representation of the cluster variables $X_k \in \mathcal{X}_q^{\mathbf{Q}}$ of the form $X_k = e^{2\pi b x_k}$ such that

- x_j is self-adjoint
- x_k satisfies the Heisenberg algebra relations

$$[x_j, x_k] = \frac{1}{2\pi i} b_{jk},$$

acting on some Hilbert space $\mathcal{H}_{\mathbf{Q}} \simeq L^2(\mathbb{R}^N)$.

Remark

Modular double \widehat{X}_k acts by $X_k^{2\pi b^{-1} x_k}$ on $\mathcal{H}_{\mathbf{Q}}$.

Polarization of $\mathcal{X}_q^{\mathbf{Q}}$

Recall $q = e^{\pi i b^2}$ such that $|q| = 1$.

Definition

A *polarization* of $\mathcal{X}_q^{\mathbf{Q}}$ is a choice of representation of the cluster variables $X_k \in \mathcal{X}_q^{\mathbf{Q}}$ of the form $X_k = e^{2\pi b x_k}$ such that

- x_j is self-adjoint
- x_k satisfies the Heisenberg algebra relations

$$[x_j, x_k] = \frac{1}{2\pi i} b_{jk},$$

acting on some Hilbert space $\mathcal{H}_{\mathbf{Q}} \simeq L^2(\mathbb{R}^N)$.

Remark

Modular double \widehat{X}_k acts by $X_k^{2\pi b^{-1}x_k}$ on $\mathcal{H}_{\mathbf{Q}}$.

Polarization of \mathcal{X}_q^Q

Example

For $X_1 X_2 = q^2 X_2 X_1$, we have

$$X_1 = e^{2\pi b x}$$

$$X_2 = e^{2\pi b p}$$

acting on $L^2(\mathbb{R})$, where $p = \frac{1}{2\pi i} \frac{d}{dx}$.

$$\textcircled{1} \mathcal{W} = \{e^{-\alpha x^2 + \beta x} \text{Poly}(x)\}$$

$\textcircled{2}$ Int. rep:

$$X_1^{is} X_2^{it} = q^{-2st} X_2^{it} X_1^{is} \quad \forall s, t \in \mathbb{R}.$$

Proposition

- Different polarizations (with the same central characters) are unitary equivalent (via $Sp(2N)$ -action)
- Cluster mutations \longleftrightarrow unitary transformation on \mathcal{H}_Q

Polarization of \mathcal{X}_q^Q

Example

For $X_1 X_2 = q^2 X_2 X_1$, we have

$$X_1 = e^{2\pi b x}$$

$$X_2 = e^{2\pi b p}$$

acting on $L^2(\mathbb{R})$, where $p = \frac{1}{2\pi i} \frac{d}{dx}$.

Proposition

- Different polarizations (with the same central characters) are unitary equivalent (via $Sp(2N)$ -action)
- Cluster mutations \longleftrightarrow unitary transformation on \mathcal{H}_Q

Polarization of \mathcal{X}_q^Q

Example

For $X_1 X_2 = q^2 X_2 X_1$, we have

$$X_1 = e^{2\pi b x}$$

$$X_2 = e^{2\pi b p}$$

acting on $L^2(\mathbb{R})$, where $p = \frac{1}{2\pi i} \frac{d}{dx}$.

Proposition


- Different polarizations (with the same central characters) are unitary equivalent (via $Sp(2N)$ -action)
- Cluster mutations \longleftrightarrow unitary transformation on \mathcal{H}_Q

Quantum cluster variety

S =Riemann surface with marked points on ∂S and punctures.

Fock-Goncharov's $\mathcal{X}_{G,S}$ -space= “(framed) local G -system”

- $\mathcal{X}_{G,S}$ has Poisson cluster \mathcal{X} variety structure \leadsto quantization $\mathcal{X}_{G,S}^q$
- To each triangle of **ideal triangulation** of S , assign a **basic quiver**.
- $G = PGL_{n+1}$: “ n -triangulation”




(Full generality: [I. (2016)], [Le (2016)], [Goncharov-Shen (2019)])

Quantum cluster variety

S =Riemann surface with marked points on ∂S and punctures.

Fock-Goncharov's $\mathcal{X}_{G,S}$ -space= “(framed) local G -system”

- $\mathcal{X}_{G,S}$ has Poisson cluster \mathcal{X} variety structure \leadsto quantization $\mathcal{X}_{G,S}^q$
- To each triangle of ideal triangulation of S , assign a basic quiver.
- $G = PGL_{n+1}$: “ n -triangulation”




(Full generality: [I. (2016)], [Le (2016)], [Goncharov-Shen (2019)])

Quantum cluster variety

S =Riemann surface with marked points on ∂S and punctures.

Fock-Goncharov's $\mathcal{X}_{G,S}$ -space= “(framed) local G -system”

- $\mathcal{X}_{G,S}$ has Poisson cluster \mathcal{X} variety structure \leadsto quantization $\mathcal{X}_{G,S}^q$
- To each triangle of **ideal triangulation** of S , assign a **basic quiver**.
- $G = PGL_{n+1}$: “ n -triangulation”




(Full generality: [I. (2016)], [Le (2016)], [Goncharov-Shen (2019)])

Quantum cluster variety

S =Riemann surface with marked points on ∂S and punctures.

Fock-Goncharov's $\mathcal{X}_{G,S}$ -space= “(framed) local G -system”

- $\mathcal{X}_{G,S}$ has Poisson cluster \mathcal{X} variety structure \leadsto quantization $\mathcal{X}_{G,S}^q$
- To each triangle of **ideal triangulation** of S , assign a **basic quiver**.
- $G = PGL_{n+1}$: “ n -triangulation”




(Full generality: [I. (2016)], [Le (2016)], [Goncharov-Shen (2019)])

Quantum cluster variety

S =Riemann surface with marked points on ∂S and punctures.

Fock-Goncharov's $\mathcal{X}_{G,S}$ -space= “(framed) local G -system”

- $\mathcal{X}_{G,S}$ has Poisson cluster \mathcal{X} variety structure \leadsto quantization $\mathcal{X}_{G,S}^q$
- To each triangle of **ideal triangulation** of S , assign a **basic quiver**.
- $G = PGL_{n+1}$: “ n -triangulation”



(Full generality: [I. (2016)], [Le (2016)], [Goncharov-Shen (2019)])

Basic Quiver

[I. (2016), Goncharov-Shen (2019)]


Definition

Elementary quiver

- $\bar{\mathbf{J}}_k(i)$, $i, k \in I$
- $Q = Q_0 = (I \setminus \{i\}) \cup \{i_l\} \cup \{i_r\} \cup \{k_e\}$

$$c_{i_l, j} = c_{j, i_r} = \frac{a_{ij}}{2}, \quad c_{i, i_r} = c_{i_r, k_e} = c_{k_e, i_l} = 1$$

- $\bar{\mathbf{J}}(i)$: without $\{k_e\}$.



Basic Quiver

[I. (2016), Goncharov-Shen (2019)]

Definition

Elementary quiver

- $\mathbf{H}(\mathbf{i})$, $\mathbf{i} = (i_1, \dots, i_m)$ reduced words
- $Q = I$

$$c_{ij} := \begin{cases} \operatorname{sgn}(r-s) \frac{a_{ij}}{2} & \beta_s = \alpha_i \text{ and } \beta_r = \alpha_j \\ 0 & \text{otherwise} \end{cases} \quad \bar{\mathbf{i}}_0: (121321)$$

- $\beta_j := s_{i_m} s_{i_{m-1}} \cdots s_{i_{j+1}}(\alpha_{i_j})$, $\alpha_i \in \Delta_+$ $\alpha_1 < \alpha_1 + d_2 < \alpha_2$
- (If $\mathbf{i} = \mathbf{i}_0$, orientation of Dynkin diagram) $< \alpha_1 + d_2 + d_3 < \alpha_2 + d_3 < \alpha_3$

$\mathbf{H}(\bar{\mathbf{i}}_0)$



Basic Quiver

[I. (2016), Goncharov-Shen (2019)]

Definition

Basic Quiver

- $\mathbf{Q}(\mathbf{i})$, $\mathbf{i} = (i_1, \dots, i_m)$ *reduced words*
- $\mathbf{Q} = \mathbf{J}_{\mathbf{i}}^{\#}(i_1) * \mathbf{J}_{\mathbf{i}}^{\#}(i_2) * \dots * \mathbf{J}_{\mathbf{i}}^{\#}(i_m) * \mathbf{H}(\mathbf{i})$
- $\mathbf{J}_{\mathbf{i}}^{\#}(i_j) = \begin{cases} \bar{\mathbf{J}}_k(i_j) & \text{if } \beta_j = \alpha_k \\ \mathbf{J}(i_j) & \text{otherwise} \end{cases}$

Basic Quiver

$$H(\tau) = \bullet$$


Example

$$\mathfrak{g} = \mathfrak{sl}_4, \mathbf{i} = (3, 2, 1).$$

$$\beta_1 = \alpha_1 + \alpha_2 + \alpha_3$$

$$\beta_2 = \alpha_1 + \alpha_2$$


$$\beta_3 = \alpha_1$$




Basic Quiver

Example

$$\mathfrak{g} = \mathfrak{sl}_4, \quad \mathbf{i}_0 = (3, 2, 1, 3, 2, 3).$$




Example: Type A_n Case



Amalgamation of 2 quivers

Example: Type A_n Case




$\mathcal{D}_{\mathfrak{sl}_{n+1}}$ -quiver $\leadsto \mathcal{X}_{\odot} := \mathcal{X}_{\mathfrak{sl}_{n+1}}$ [Schrader-Shapiro]

$$\iota : \mathcal{D}_q(\mathfrak{sl}_{n+1}) \hookrightarrow \mathcal{X}_{\odot}$$

$$\mathcal{U}_q(\mathfrak{sl}_{n+1}) \hookrightarrow \mathcal{X}_{\odot} / \langle \iota(K_i) \iota(K'_i) = 1 \rangle$$

Example: Type A_n Case



Embedding of $F_i \in \mathfrak{D}_{\mathfrak{sl}_4} \hookrightarrow \mathcal{X}_\odot$

$$f_1 = X_1 + X_{1,2} + X_{1,2,3} + X_{1,2,3,4} + X_{1,2,3,4,5} + X_{1,2,3,4,5,6}$$


$$f_2 = X_8 + X_{8,9} + X_{8,9,10} + X_{8,9,10,11}$$

$$f_3 = X_{13} + X_{13,14}$$

$$K'_1 = X_{1,2,3,4,5,6,7} \quad K'_2 = X_{8,9,10,11,12} \quad K'_3 = X_{13,14,15}$$

$$\in \mathbb{C}[q][x_i]$$

Example: Type A_n Case



Embedding of $E_i \in \mathfrak{D}_{\mathfrak{sl}_4} \hookrightarrow \mathcal{X}_\odot$

$$e_1 = X_7 + X_{7,16}$$

$$e_2 = X_{12} + X_{12,6} + X_{12,6,17} + X_{12,6,17,2}$$

$$e_3 = X_{15} + X_{15,11} + X_{15,11,5} + X_{15,11,5,18} + X_{15,11,5,18,3} + X_{15,11,5,18,3,9}$$

$$K_1 = X_{7,16,1} \quad K_2 = X_{12,6,17,2,8} \quad K_3 = X_{15,11,5,18,3,9,13}$$

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Theorem (Schrader-Shapiro, I. (2016))

- *There exists an embedding*

$$\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_{\odot}$$

corresponding to the quiver $\mathcal{D}_{\mathfrak{g}}$ associated to



- *We recover the positive representations $\mathcal{P}_{\lambda} \simeq \mathcal{H}_{\odot}$ through a polarization of \mathcal{X}_{\odot} .*

Theorem (I. (2016))

- *The generators $\mathbf{e}_i, \mathbf{f}_i, K_i$ are represented by **positive polynomials** (i.e. over $\mathbb{N}[q, q^{-1}]$) in the cluster variables $X_i \in \mathcal{X}_{\odot}$.*
- *The generators $\mathbf{e}_i, \mathbf{f}_i, K_i$ are labeled by paths on the $\mathcal{D}_{\mathfrak{g}}$ quiver.*
- *\mathbf{f}_i paths are simple - coincide with **Feigin's homomorphism**.*


Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Theorem (Schrader-Shapiro, I. (2016))

- There exists an embedding

$$\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_{\odot}$$

g general type

corresponding to the quiver $\mathcal{D}_{\mathfrak{g}}$ associated to 

- We recover the positive representations $\mathcal{P}_{\lambda} \simeq \mathcal{H}_{\odot}$ through a polarization of \mathcal{X}_{\odot} .

Theorem (I. (2016))

- The generators $\mathbf{e}_i, \mathbf{f}_i, K_i$ are represented by *positive polynomials* (i.e. over $\mathbb{N}[q, q^{-1}]$) in the cluster variables $X_i \in \mathcal{X}_{\odot}$.
- The generators $\mathbf{e}_i, \mathbf{f}_i, K_i$ are labeled by paths on the $\mathcal{D}_{\mathfrak{g}}$ quiver.
- \mathbf{f}_i paths are simple - coincide with *Feigin's homomorphism*.

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Theorem (Schrader-Shapiro, I. (2016))

- *There exists an embedding*

$$\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_{\odot}$$

corresponding to the quiver $\mathcal{D}_{\mathfrak{g}}$ associated to



- *We recover the positive representations $\mathcal{P}_{\lambda} \simeq \mathcal{H}_{\odot}$ through a polarization of \mathcal{X}_{\odot} .*

Theorem (I. (2016))

- *The generators $\mathbf{e}_i, \mathbf{f}_i, K_i$ are represented by **positive polynomials** (i.e. over $\mathbb{N}[q, q^{-1}]$) in the cluster variables $X_i \in \mathcal{X}_{\odot}$.*
- *The generators $\mathbf{e}_i, \mathbf{f}_i, K_i$ are labeled by paths on the $\mathcal{D}_{\mathfrak{g}}$ quiver.*
- *\mathbf{f}_i paths are simple - coincide with **Feigin's homomorphism**.*


Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Theorem (Schrader-Shapiro, I. (2016))

- *There exists an embedding*

$$\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_{\odot}$$

corresponding to the quiver $\mathcal{D}_{\mathfrak{g}}$ associated to



- *We recover the positive representations $\mathcal{P}_{\lambda} \simeq \mathcal{H}_{\odot}$ through a polarization of \mathcal{X}_{\odot} .*

Theorem (I. (2016))


- *The generators $\mathbf{e}_i, \mathbf{f}_i, K_i$ are represented by **positive polynomials** (i.e. over $\mathbb{N}[q, q^{-1}]$) in the cluster variables $X_i \in \mathcal{X}_{\odot}$.*
- *The generators $\mathbf{e}_i, \mathbf{f}_i, K_i$ are labeled by paths on the $\mathcal{D}_{\mathfrak{g}}$ quiver.*
- *\mathbf{f}_i paths are simple - coincide with **Feigin's homomorphism**.*

Positive Representations of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$

Theorem (Schrader-Shapiro, I. (2016))


- There exists an embedding

$$\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_{\odot}$$

corresponding to the quiver $\mathcal{D}_{\mathfrak{g}}$ associated to 


- We recover the positive representations $\mathcal{P}_{\lambda} \simeq \mathcal{H}_{\odot}$ through a polarization of \mathcal{X}_{\odot} .

Theorem (I. (2016))

- The generators $\mathbf{e}_i, \mathbf{f}_i, K_i$ are represented by *positive polynomials* (i.e. over $\mathbb{N}[q, q^{-1}]$) in the cluster variables $X_i \in \mathcal{X}_{\odot}$.
- The generators $\mathbf{e}_i, \mathbf{f}_i, K_i$ are labeled by paths on the $\mathcal{D}_{\mathfrak{g}}$ quiver.
- \mathbf{f}_i paths are simple - coincide with *Feigin's homomorphism*. 


E_6 embedding

$$\mathbf{i}_0 = (3 \ 43 \ 034 \ 230432 \ 12340321 \ 5432103243054321)$$



E_6 embedding

$$\mathbf{i}_0 = (3 \ 43 \ 034 \ 230432 \ 12340321 \ 5432103243054321)$$



Minimal Positive Representation for $\mathcal{U}_q(\mathfrak{sl}(n+1, \mathbb{R}))$

Minimal Positive Representation

- Parabolic subgroups $\longleftrightarrow J \subset I$
- $P_J := B_- L_J$, Levi subgroup $L_J = \langle T, U_j^+, U_j^- \rangle_{j \in J}$
- $P_\emptyset := B_-$.

Example

For $G = SL_4$, $J = \{1, 2\} \subset I = \{1, 2, 3\}$

$$P_J = \begin{pmatrix} * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix}$$

$$(G/P_J)_{>0} = \begin{pmatrix} * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix} \begin{pmatrix} 1 & a & 0 & 0 \\ 0 & 1 & b & 0 \\ 0 & 0 & 1 & c \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad a, b, c > 0$$

$$x_3(c)x_2(b)x_1(a)e^{tX} = n \cdot h \cdot x_1(f')x_2(e')x_1(d')x_3(c')x_2(b')x_1(a'), \quad n \in U_-, h \in T$$

Minimal Positive Representation

- Parabolic subgroups $\longleftrightarrow J \subset I$
- $P_J := B_- L_J$, Levi subgroup $L_J = \langle T, U_j^+, U_j^- \rangle_{j \in J}$
- $P_\emptyset := B_-$.

Example

For $G = SL_4$, $J = \{1, 2\} \subset I = \{1, 2, 3\}$

$$P_J = \begin{pmatrix} * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix}$$

$$(G/P_J)_{>0} = \begin{pmatrix} * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix} \begin{pmatrix} 1 & a & 0 & 0 \\ 0 & 1 & b & 0 \\ 0 & 0 & 1 & c \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad a, b, c > 0$$

$$x_3(c)x_2(b)x_1(a)e^{tX} = n \cdot h \cdot x_1(f')x_2(e')x_1(d')x_3(c')x_2(b')x_1(a'), \quad n \in U_-, h \in T$$

Minimal Positive Representation

- Parabolic subgroups $\longleftrightarrow J \subset I$
- $P_J := B_- L_J$, Levi subgroup $L_J = \langle T, U_j^+, U_j^- \rangle_{j \in J}$
- $P_\emptyset := B_-$.

Example

For $G = SL_4$, $J = \{1, 2\} \subset I = \{1, 2, 3\}$

$$P_J = \begin{pmatrix} * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix}$$

$$(G/P_J)_{>0} = \begin{pmatrix} * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix} \begin{pmatrix} 1 & a & 0 & 0 \\ 0 & 1 & b & 0 \\ 0 & 0 & 1 & c \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad a, b, c > 0$$

$$x_3(c)x_2(b)x_1(a)e^{tX} = n \cdot h \cdot x_1(f')x_2(e')x_1(d')x_3(c')x_2(b')x_1(a'), \quad n \in U_-, h \in T$$

Minimal Positive Representation

- Parabolic subgroups $\longleftrightarrow J \subset I$
- $P_J := B_- L_J$, Levi subgroup $L_J = \langle T, U_j^+, U_j^- \rangle_{j \in J}$
- $P_\emptyset := B_-$.

Example

For $G = SL_4$, $J = \{1, 2\} \subset I = \{1, 2, 3\}$

$$P_J = \begin{pmatrix} * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix}$$

$$(G/P_J)_{>0} = \begin{pmatrix} * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix} \begin{pmatrix} 1 & a & 0 & 0 \\ 0 & 1 & b & 0 \\ 0 & 0 & 1 & c \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad a, b, c > 0$$

$$x_3(c)x_2(b)x_1(a)e^{tX} = n \cdot h \cdot x_1(f')x_2(e')x_1(d')x_3(c')x_2(b')x_1(a'), \quad n \in U_-, h \in T$$

Minimal Positive Representation

- Parabolic subgroups $\longleftrightarrow J \subset I$
- $P_J := B_- L_J$, Levi subgroup $L_J = \langle T, U_j^+, U_j^- \rangle_{j \in J}$
- $P_\emptyset := B_-$.

Example

For $G = SL_4$, $J = \{1, 2\} \subset I = \{1, 2, 3\}$

$$P_J = \begin{pmatrix} * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix}$$

$$(G/P_J)_{>0} = \begin{pmatrix} * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix} \begin{pmatrix} 1 & a & 0 & 0 \\ 0 & 1 & b & 0 \\ 0 & 0 & 1 & c \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad a, b, c > 0$$

$$x_3(c)x_2(b)x_1(a)e^{tX} = n \cdot h \cdot x_1(f')x_2(e')x_1(d')x_3(c')x_2(b')x_1(a'), \quad n \in U_-, h \in T$$

Minimal Positive Representation

- Parabolic subgroups $\longleftrightarrow J \subset I$
- $P_J := B_- L_J$, Levi subgroup $L_J = \langle T, U_j^+, U_j^- \rangle_{j \in J}$
- $P_\emptyset := B_-$.

Example

For $G = SL_4$, $J = \{1, 2\} \subset I = \{1, 2, 3\}$

$$P_J = \begin{pmatrix} * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix}$$

$$(G/P_J)_{>0} = \begin{pmatrix} * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix} \begin{pmatrix} 1 & a & 0 & 0 \\ 0 & 1 & b & 0 \\ 0 & 0 & 1 & c \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad a, b, c > 0$$

$$x_3(c)x_2(b)x_1(a)e^{tX} = n \cdot h \cdot x_1(f')x_2(e')x_1(d')x_3(c')x_2(b')x_1(a'), \quad n \in U_-, h \in T$$

Minimal Positive Representation

- Parabolic subgroups $\longleftrightarrow J \subset I$
- $P_J := B_- L_J$, Levi subgroup $L_J = \langle T, U_j^+, U_j^- \rangle_{j \in J}$
- $P_\emptyset := B_-$.

Example

For $G = SL_4$, $J = \{1, 2\} \subset I = \{1, 2, 3\}$

$$P_J = \begin{pmatrix} * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix}$$

$$(G/P_J)_{>0} = \begin{pmatrix} * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & 0 \\ * & * & * & * \end{pmatrix} \begin{pmatrix} 1 & a & 0 & 0 \\ 0 & 1 & b & 0 \\ 0 & 0 & 1 & c \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad a, b, c > 0$$

$$x_3(c)x_2(b)x_1(a)e^{tX} = \underbrace{h}_{\text{circled}} \underbrace{x_1(a')x_2(b')x_3(c')}_{\text{crossed out}}, \quad n \in U_-, h \in T$$

Minimal Positive Representation

Previous recipe produces a representation \mathcal{P}_λ^J for $\mathcal{U}_q(\mathfrak{sl}(4, \mathbb{R}))$, ($\lambda \in \mathbb{R}$)

$$\pi_\lambda^J(\mathbf{e}_1) = e^{\pi b(u-2p_u)} + e^{\pi b(-u-2p_u)}$$

$$\pi_\lambda^J(\mathbf{e}_2) = e^{\pi b(-u+v-2p_v)} + e^{\pi b(u-v-2p_v)}$$

$$\pi_\lambda^J(\mathbf{e}_3) = e^{\pi b(-v+w-2p_w)} + e^{\pi b(v-w-2p_w)}$$

$$\pi_\lambda^J(\mathbf{f}_1) = e^{\pi b(-u+v+2p_u)} + e^{\pi b(u-v+2p_u)}$$

$$\pi_\lambda^J(\mathbf{f}_2) = e^{\pi b(-v+w+2p_v)} + e^{\pi b(v-w+2p_v)}$$

$$\pi_\lambda^J(\mathbf{f}_3) = e^{\pi b(2\lambda-w+2p_w)} + e^{\pi b(-2\lambda+w+2p_w)}$$

$$\pi_\lambda^J(\mathbf{K}_1) = e^{\pi b(-2u+v)}$$


$$\pi_\lambda^J(\mathbf{K}_2) = e^{\pi b(u-2v+w)}$$

$$\pi_\lambda^J(\mathbf{K}_3) = e^{\pi b(v-2w+2\lambda)}$$

Xi

acting on $L^2(\mathbb{R}^3)$ as positive self-adjoint operators.

Minimal Positive Representation



w_0 I
 w_J J

$w_J^{-1} w_0$

$$D(i) := Q(i^{op}) * Q(i), \quad i = (3, 2, 1)$$

$$e_1 = X_3 + X_{3,0}$$

$$e_2 = X_6 + X_{6,2}$$

$$e_3 = X_9 + X_{9,5}$$

$$f_1 = X_1 + X_{1,2}$$

$$f_2 = X_4 + X_{4,5}$$

$$f_3 = X_7 + X_{7,8}$$

$$K_1 = X_{3,0,1}$$

$$K_2 = X_{6,2,4}$$

$$K_3 = X_{9,5,7}$$

$$K'_1 = X_{1,2,3}$$

$$K'_2 = X_{4,5,6}$$


$$K'_3 = X_{7,8,9}$$

$$\text{Central character: } \pi(X_{0,2,5,8}) = e^{-4\pi b\lambda}$$

Minimal Positive Representation

Theorem (I. (2020))

The polarization of the quiver $\mathbf{D}(\mathbf{i})$ for $\mathbf{i} = (n, \dots, 3, 2, 1)$ gives a representation \mathcal{P}_λ^J of $\mathcal{U}_q(\mathfrak{sl}(n+1, \mathbb{R}))$ acting on $L^2(\mathbb{R}^n)$ as positive self-adjoint operators.



Minimal Positive Representation

Theorem (I. (2020))

- *The non-simple generators*

$$\mathbf{e}_\alpha := T_{i_1} \cdots T_{i_{k-1}}(\mathbf{e}_k) \quad \text{pos. self ad.}$$

$$\mathbf{f}_\alpha := T_{i_1} \cdots T_{i_{k-1}}(\mathbf{f}_k)$$

is non-zero, where T_i = Lusztig's braid group action.

- *The universal \mathcal{R} operator is well-defined*

$$\mathcal{R} = \mathcal{K} \prod_{\alpha \in \Phi_+} g_\alpha(\mathbf{e}_\alpha \otimes \mathbf{f}_\alpha)$$

- *The Casimirs \mathbf{C}_k acts by real-valued scalar, and lie outside the positive spectrum of the usual positive representations.*

Minimal Positive Representation

Theorem (I. (2020))

- *The non-simple generators*

$$\mathbf{e}_\alpha := T_{i_1} \cdots T_{i_{k-1}}(\mathbf{e}_k)$$

$$\mathbf{f}_\alpha := T_{i_1} \cdots T_{i_{k-1}}(\mathbf{f}_k)$$

is non-zero, where T_i = Lusztig's braid group action.

- *The universal \mathcal{R} operator is well-defined*

$$\mathcal{R} = \mathcal{K} \prod_{\alpha \in \Phi_+} g_\alpha(\mathbf{e}_\alpha \otimes \mathbf{f}_\alpha)$$

↑ quantum. diag.

- *The Casimirs \mathbf{C}_k acts by real-valued scalar, and lie outside the positive spectrum of the usual positive representations.*

Minimal Positive Representation

Theorem (I. (2020))

- *The non-simple generators*

$$\mathbf{e}_\alpha := T_{i_1} \cdots T_{i_{k-1}}(\mathbf{e}_k)$$

$$\mathbf{f}_\alpha := T_{i_1} \cdots T_{i_{k-1}}(\mathbf{f}_k)$$

is non-zero, where T_i = Lusztig's braid group action.

- *The universal \mathcal{R} operator is well-defined*


$$\mathcal{R} = \mathcal{K} \prod_{\alpha \in \Phi_+} g_\alpha(\mathbf{e}_\alpha \otimes \mathbf{f}_\alpha)$$

- *The Casimirs \mathbf{C}_k acts by real-valued scalar, and lie outside the [positive spectrum](#) of the usual positive representations.*


Casimirs

Example


$\mathcal{U}_q(\mathfrak{sl}(3, \mathbb{R}))$, the possible action of (C_1, C_2) (by scalars) on \mathcal{P}_λ and \mathcal{P}_λ^J :




Evaluation Module of $\mathcal{U}_q(\widehat{\mathfrak{sl}}_{n+1})$



Evaluation Module of $\mathcal{U}_q(\widehat{\mathfrak{sl}}_{n+1})$



Evaluation Module of $\mathcal{U}_q(\widehat{\mathfrak{sl}}_{n+1})$



Evaluation Module of $\mathcal{U}_q(\widehat{\mathfrak{sl}}_{n+1})$

Theorem (I. (2020))

The positive representation of $\mathcal{U}_q(\widehat{\mathfrak{sl}}_{n+1})$ defined by the polarization of the previous quiver is unitarily equivalent to Jimbo's evaluation module \mathcal{P}_λ^μ , $\mu \in \mathbb{R}$

$$\overset{\textcolor{red}{\curvearrowright}}{\mathcal{E}} \mapsto \mathcal{E}$$

$$\mathcal{U}_q(\widehat{\mathfrak{sl}}_{n+1}) \longrightarrow \mathcal{U}_q(\mathfrak{sl}_{n+1})$$


of the minimal positive representations \mathcal{P}_λ^J of $\mathcal{U}_q(\mathfrak{sl}_{n+1})$, where

$$e^{\pi b \mu} := \pi(D_0^{\frac{1}{n+1}} D_1)$$

(D_0 =product of all middle vertices, D_1 = product of all right vertices.)

Positive representation of $\mathcal{U}_q(\widehat{\mathfrak{sl}}_2)$

Example



$$\mathbf{f}_0 = X_1 + X_{1,2}$$

$$\mathbf{e}_0 = X_3 + X_{3,5}$$

$$\mathbf{f}_1 = X_4 + X_{4,5}$$

$$\mathbf{e}_1 = X_6 + X_{6,2}$$

Serre relation ($a_{01} = a_{10} = -2$):

$$X_i^3 X_j - [3]_q X_i^2 X_j X_i + [3]_q X_i X_j X_i^2 - X_j X_i^3 = 0, \quad i \neq j$$

General Construction

Main Theorem

Parabolic induction \longleftrightarrow truncating $\mathbf{i}_J \subset \mathbf{i}_0$ where $\mathbf{i}_J, \mathbf{i}_0$ are the longest word of the Weyl groups $W_J \subset W$.

$$w_0 = w_J \bar{w}$$

$$\bar{w} \longleftrightarrow \bar{\mathbf{i}}$$

Example

$$W_{\mathfrak{sl}_4} \subset W_{\mathfrak{sl}_5}$$

$$\mathbf{i}_0 = (1, 2, 1, 3, 2, 1, 4, 3, 2, 1)$$

Observe that

$$\mathbf{Q}(\mathbf{i}) = \mathbf{Q}(\mathbf{i}_J) * \mathbf{Q}(\bar{\mathbf{i}})$$

In general, we have realization of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ on the quantum torus algebra associated to the symplectic double $\mathbf{D}(\mathbf{i})$.

Main Theorem

Parabolic induction \longleftrightarrow truncating $\mathbf{i}_J \subset \mathbf{i}_0$ where $\mathbf{i}_J, \mathbf{i}_0$ are the longest word of the Weyl groups $W_J \subset W$.

$$w_0 = w_J \bar{w}$$

$$\bar{w} \longleftrightarrow \bar{\mathbf{i}}$$

Example

$$W_{\mathfrak{sl}_4} \subset W_{\mathfrak{sl}_5}$$

$$\mathbf{i}_0 = (1, 2, 1, 3, 2, 1, 4, 3, 2, 1)$$

Observe that

$$\mathbf{Q}(\mathbf{i}) = \mathbf{Q}(\mathbf{i}_J) * \mathbf{Q}(\bar{\mathbf{i}})$$

In general, we have realization of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ on the quantum torus algebra associated to the symplectic double $\mathbf{D}(\mathbf{i})$.

Main Theorem

Parabolic induction \longleftrightarrow truncating $\mathbf{i}_J \subset \mathbf{i}_0$ where $\mathbf{i}_J, \mathbf{i}_0$ are the longest word of the Weyl groups $W_J \subset W$.

$$w_0 = w_J \bar{w}$$

$$\bar{w} \longleftrightarrow \bar{\mathbf{i}}$$

Example

$$W_{\mathfrak{sl}_4} \subset W_{\mathfrak{sl}_5}$$

$$\mathbf{i}_0 = (1, 2, 1, 3, 2, 1, 4, 3, 2, 1)$$

\mathbf{i}_J
 $\bar{\mathbf{i}}$

Observe that

$$\mathbf{Q}(\mathbf{i}_0) = \mathbf{Q}(\mathbf{i}_J) * \mathbf{Q}(\bar{\mathbf{i}})$$

In general, we have realization of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ on the quantum torus algebra associated to the symplectic double $\mathbf{D}(\bar{\mathbf{i}})$.

Main Theorem

Theorem (I. (2020))

- There is a homomorphism

$$\mathcal{D}_q(\mathfrak{g}) \longrightarrow \mathcal{X}_q^{\mathbf{D}(\bar{\mathfrak{i}})}$$

\swarrow $\mathbb{Q}^{\text{op}} * \mathbb{Q}$

such that the image of universally Laurent polynomials.

- A polarization of $\mathcal{X}_q^{\mathbf{D}(\bar{\mathfrak{i}})}$ induces a family of irreducible representations \mathcal{P}_λ^J of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ parametrized by $\lambda \in \mathbb{R}^{I \setminus J}$ as positive self-adjoint operators on $L^2(\mathbb{R}^{I(\bar{w})})$.

Corollary

The *parabolic positive representations* \mathcal{P}_λ^J is obtained as a quantum twist of the parabolic induction, by ignoring the variables u_i corresponding to the Levi subgroups L_J of P_J in the quotient G/P_J .

\iff setting formally $e^{\pi b u_i} = 1$ and $e^{\pm \pi b p_i} = 0$.

Main Theorem

Theorem (I. (2020))

- *There is a homomorphism*

$$\mathcal{D}_q(\mathfrak{g}) \longrightarrow \mathcal{X}_q^{\mathbf{D}(\bar{\mathfrak{i}})}$$

such that the image of universally Laurent polynomials.

- *A polarization of $\mathcal{X}_q^{\mathbf{D}(\bar{\mathfrak{i}})}$ induces a family of irreducible representations \mathcal{P}_λ^J of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ parametrized by $\lambda \in \mathbb{R}^{I \setminus J}$ as positive self-adjoint operators on $L^2(\mathbb{R}^{l(\bar{w})})$.*

Corollary

The parabolic positive representations \mathcal{P}_λ^J is obtained as a quantum twist of the parabolic induction, by ignoring the variables u_i corresponding to the Levi subgroups L_J of P_J in the quotient G/P_J .

\iff *setting formally $e^{\pi b u_i} = 1$ and $e^{\pm \pi b p_i} = 0$.*

Main Theorem

Theorem (I. (2020))

- *There is a homomorphism*

$$\mathcal{D}_q(\mathfrak{g}) \longrightarrow \mathcal{X}_q^{\mathbf{D}(\bar{\mathfrak{i}})}$$

such that the image of universally Laurent polynomials.

- *A polarization of $\mathcal{X}_q^{\mathbf{D}(\bar{\mathfrak{i}})}$ induces a family of irreducible representations \mathcal{P}_λ^J of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ parametrized by $\lambda \in \mathbb{R}^{I \setminus J}$ as positive self-adjoint operators on $L^2(\mathbb{R}^{l(\bar{w})})$.*

Corollary

*The **parabolic positive representations** \mathcal{P}_λ^J is obtained as a quantum twist of the parabolic induction, by ignoring the variables u_i corresponding to the Levi subgroups L_J of P_J in the quotient G/P_J .*

\iff setting formally $e^{\pi b u_i} = 1$ and $e^{\pm \pi b p_i} = 0$.

Main Theorem

Theorem (I. (2020))

- *There is a homomorphism*

$$\mathcal{D}_q(\mathfrak{g}) \longrightarrow \mathcal{X}_q^{\mathbf{D}(\bar{\mathfrak{i}})}$$

such that the image of universally Laurent polynomials.

- *A polarization of $\mathcal{X}_q^{\mathbf{D}(\bar{\mathfrak{i}})}$ induces a family of irreducible representations \mathcal{P}_λ^J of $\mathcal{U}_q(\mathfrak{g}_{\mathbb{R}})$ parametrized by $\lambda \in \mathbb{R}^{I \setminus J}$ as positive self-adjoint operators on $L^2(\mathbb{R}^{l(\bar{w})})$.*

Corollary

*The **parabolic positive representations** \mathcal{P}_λ^J is obtained as a quantum twist of the parabolic induction, by ignoring the variables u_i corresponding to the Levi subgroups L_J of P_J in the quotient G/P_J .*

\iff *setting formally $e^{\pi b u_i} = 1$ and $e^{\pm \pi b p_i} = 0$.*

Idea of Proof

Definition

The *Heisenberg double* $\mathcal{H}_q^\pm(\mathfrak{g}) := \langle \mathbf{e}_i^\pm, \mathbf{f}_i^\pm, \mathbf{K}_i^\pm, \mathbf{K}'_i{}^\pm \rangle$ satisfying

$$\frac{[\mathbf{e}_i^+, \mathbf{f}_j^+]}{q - q^{-1}} = \delta_{ij} \mathbf{K}'_i{}^+, \quad \frac{[\mathbf{e}_i^-, \mathbf{f}_j^-]}{q - q^{-1}} = \delta_{ij} \mathbf{K}_i^-$$

and other standard quantum group relations.

Proposition

The embedding $\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{\mathbf{D}(\mathbf{i}_0)} \subset \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_0^{op})} \otimes \mathcal{X}_q^{\mathbf{Q}(\mathbf{i})}$ decomposes as

$$\begin{aligned} \mathbf{e}_i &= \mathbf{e}_i^+ + \mathbf{K}_i^+ \mathbf{e}_i^-, & \mathbf{f}_i &= \mathbf{f}_i^- + \mathbf{K}'_i{}^- \mathbf{f}_i^+ \\ \mathbf{K}_i &= \mathbf{K}_i^+ \mathbf{K}_i^-, & \mathbf{K}'_i &= \mathbf{K}'_i{}^+ \mathbf{K}'_i{}^- \end{aligned}$$

where $\mathcal{H}_q^+(\mathfrak{g}) \hookrightarrow 1 \otimes \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_0)}$, $\mathcal{H}_q^-(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_0^{op})} \otimes 1$

Idea of Proof

Definition

The *Heisenberg double* $\mathcal{H}_q^\pm(\mathfrak{g}) := \langle \mathbf{e}_i^\pm, \mathbf{f}_i^\pm, \mathbf{K}_i^\pm, \mathbf{K}'_i{}^\pm \rangle$ satisfying

$$\frac{[\mathbf{e}_i^+, \mathbf{f}_j^+]}{q - q^{-1}} = \delta_{ij} \mathbf{K}'_i{}^+, \quad \frac{[\mathbf{e}_i^-, \mathbf{f}_j^-]}{q - q^{-1}} = \delta_{ij} \mathbf{K}_i^-$$

and other standard quantum group relations.

Proposition

The embedding $\mathcal{D}_q(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{\mathbf{D}(\mathbf{i}_0)} \subset \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_0^{op})} \otimes \mathcal{X}_q^{\mathbf{Q}(\mathbf{i})}$ decomposes as

$$\begin{aligned} \mathbf{e}_i &= \mathbf{e}_i^+ + \mathbf{K}_i^+ \mathbf{e}_i^-, & \mathbf{f}_i &= \mathbf{f}_i^- + \mathbf{K}'_i{}^- \mathbf{f}_i^+ \\ \mathbf{K}_i &= \mathbf{K}_i^+ \mathbf{K}_i^-, & \mathbf{K}'_i &= \mathbf{K}'_i{}^+ \mathbf{K}'_i{}^- \end{aligned}$$

where $\mathcal{H}_q^+(\mathfrak{g}) \hookrightarrow 1 \otimes \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_0)}$, $\mathcal{H}_q^-(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_0^{op})} \otimes 1$



Idea of Proof

Definition

The *generalized Heisenberg double* $\mathcal{H}_{q,\omega}^{\pm}(\mathfrak{g}) := \langle \mathbf{e}_i^{\pm}, \mathbf{f}_i^{\pm}, \mathbf{K}_i^{\pm}, \mathbf{K}'_i^{\pm} \rangle$

$$\frac{[\mathbf{e}_i^+, \mathbf{f}_j^+]}{q - q^{-1}} = \delta_{ij} \mathbf{K}'_i^+ + \omega_{ij} \mathbf{K}_i^+, \quad \frac{[\mathbf{e}_i^-, \mathbf{f}_j^-]}{q - q^{-1}} = \delta_{ij} \mathbf{K}_i^- - \omega_{ij} \mathbf{K}'_i^-$$

and other standard quantum group relations, where $\omega_{ij} \in \mathbb{C}$.

Proposition

If $\mathcal{H}_{q,\omega}^{\pm}(\mathfrak{g})$ are commuting copies, then

$$\begin{aligned} \mathbf{e}_i &= \mathbf{e}_i^+ + \mathbf{K}_i^+ \mathbf{e}_i^-, & \mathbf{f}_i &= \mathbf{f}_i^- + \mathbf{K}_i'^- \mathbf{f}_i^+ \\ \mathbf{K}_i &= \mathbf{K}_i^+ \mathbf{K}_i^-, & \mathbf{K}'_i &= \mathbf{K}'_i^+ \mathbf{K}'_i^- \end{aligned}$$

gives a homomorphic image of $\mathcal{U}_q(\mathfrak{g})$.

Idea of Proof

Definition


Let $J \subset I$. The *double Dynkin involution* of $i \in I$ is the unique index $i^{**} \in I$ such that

$$w_0 s_i = s_{i^*} w_0 = s_{i^*} w_J \bar{w} = w_J s_{i^{**}} \bar{w}.$$

$$\iff i^{**} := (i^{*w})^{*w_J}$$

where $i^{*w_J} = i$ if $i \notin J$.

Ex
 $D_5 \subset E_6$



$$2^{**J} = 4^{*J} = 0$$

Decomposition Lemma



Lemma (*Decomposition Lemma*)

The embedding $\mathcal{H}_q^+(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_0)} \subset \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_J)} \otimes \mathcal{X}_q^{\mathbf{Q}(\bar{\mathbf{i}})}$ can be decomposed into the form

$$\begin{aligned} e_i^+ &= \overline{e_i} + \overline{K_i} e_{i**}^J, & f_i^+ &= f_i^J + K_i'^J \overline{f_i}, \\ K_i^+ &= K_{i**}^J \overline{K_i}, & K_i'^+ &= K_i'^J \overline{K_i'} \end{aligned}$$

where $e_i^J = f_i^J = 0$ and $K_i^J = K_i'^J = 1$ if $i \notin J$, such that

- $X_i^J \in \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_J)} \otimes 1$ and $\overline{X_i} \in 1 \otimes \mathcal{X}_q^{\mathbf{Q}(\bar{\mathbf{i}})}$ for $X = e, f, K, K'$
- $\{e_i^J, f_i^J, K_i^J, K_i'^J\} \simeq \mathcal{H}_q^+(\mathfrak{g}_J)$ in $\mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_J)}$ where $\mathfrak{g}_J \subset \mathfrak{g}$.
- We have on $\mathcal{X}_q^{\mathbf{Q}(\bar{\mathbf{i}})}$ for some $\omega_{ij} \in \{0, 1\}$,

$$\langle \overline{e_i}, \overline{f_i}, \overline{K_i}, \overline{K_i'} \rangle \simeq \mathcal{H}_{q,\omega}^+(\mathfrak{g})$$

Decomposition Lemma

Lemma (*Decomposition Lemma*)

The embedding $\mathcal{H}_q^+(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_0)} \subset \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_J)} \otimes \mathcal{X}_q^{\mathbf{Q}(\bar{\mathbf{i}})}$ can be decomposed into the form

$$\begin{aligned} e_i^+ &= \overline{e_i} + \overline{K_i} e_{i**}^J, & f_i^+ &= f_i^J + K_i'^J \overline{f_i}, \\ K_i^+ &= K_{i**}^J \overline{K_i}, & K_i'^+ &= K_i'^J \overline{K_i'} \end{aligned}$$

where $e_i^J = f_i^J = 0$ and $K_i^J = K_i'^J = 1$ if $i \notin J$, such that

- $X_i^J \in \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_J)} \otimes 1$ and $\overline{X_i} \in 1 \otimes \mathcal{X}_q^{\mathbf{Q}(\bar{\mathbf{i}})}$ for $X = e, f, K, K'$
- $\{e_i^J, f_i^J, K_i^J, K_i'^J\} \simeq \mathcal{H}_q^+(\mathfrak{g}_J)$ in $\mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_J)}$ where $\mathfrak{g}_J \subset \mathfrak{g}$.
- We have on $\mathcal{X}_q^{\mathbf{Q}(\bar{\mathbf{i}})}$ for some $\omega_{ij} \in \{0, 1\}$,

$$\langle \overline{e_i}, \overline{f_i}, \overline{K_i}, \overline{K_i'} \rangle \simeq \mathcal{H}_{q,\omega}^+(\mathfrak{g})$$

Decomposition Lemma

Lemma (*Decomposition Lemma*)

The embedding $\mathcal{H}_q^+(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_0)} \subset \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_J)} \otimes \mathcal{X}_q^{\mathbf{Q}(\bar{\mathbf{i}})}$ can be decomposed into the form

$$\begin{aligned} e_i^+ &= \overline{e_i} + \overline{K_i} e_{i**}^J, & f_i^+ &= f_i^J + K_i'^J \overline{f_i}, \\ K_i^+ &= K_{i**}^J \overline{K_i}, & K_i'^+ &= K_i'^J \overline{K_i'} \end{aligned}$$

where $e_i^J = f_i^J = 0$ and $K_i^J = K_i'^J = 1$ if $i \notin J$, such that

- $X_i^J \in \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_J)} \otimes 1$ and $\overline{X_i} \in 1 \otimes \mathcal{X}_q^{\mathbf{Q}(\bar{\mathbf{i}})}$ for $X = e, f, K, K'$
- $\{e_i^J, f_i^J, K_i^J, K_i'^J\} \simeq \mathcal{H}_q^+(\mathfrak{g}_J)$ in $\mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_J)}$ where $\mathfrak{g}_J \subset \mathfrak{g}$.
- We have on $\mathcal{X}_q^{\mathbf{Q}(\bar{\mathbf{i}})}$ for some $\omega_{ij} \in \{0, 1\}$,

$$\langle \overline{e_i}, \overline{f_i}, \overline{K_i}, \overline{K_i'} \rangle \simeq \mathcal{H}_{q,\omega}^+(\mathfrak{g})$$

Decomposition Lemma

Lemma (*Decomposition Lemma*)

The embedding $\mathcal{H}_q^+(\mathfrak{g}) \hookrightarrow \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_0)} \subset \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_J)} \otimes \mathcal{X}_q^{\mathbf{Q}(\bar{\mathbf{i}})}$ can be decomposed into the form

$$\begin{aligned} e_i^+ &= \overline{e_i} + \overline{K_i} e_{i**}^J, & f_i^+ &= f_i^J + K_i'^J \overline{f_i}, \\ K_i^+ &= K_{i**}^J \overline{K_i}, & K_i'^+ &= K_i'^J \overline{K_i'} \end{aligned}$$

where $e_i^J = f_i^J = 0$ and $K_i^J = K_i'^J = 1$ if $i \notin J$, such that

- $X_i^J \in \mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_J)} \otimes 1$ and $\overline{X_i} \in 1 \otimes \mathcal{X}_q^{\mathbf{Q}(\bar{\mathbf{i}})}$ for $X = e, f, K, K'$
- $\{e_i^J, f_i^J, K_i^J, K_i'^J\} \simeq \mathcal{H}_q^+(\mathfrak{g}_J)$ in $\mathcal{X}_q^{\mathbf{Q}(\mathbf{i}_J)}$ where $\mathfrak{g}_J \subset \mathfrak{g}$.
- We have on $\mathcal{X}_q^{\mathbf{Q}(\bar{\mathbf{i}})}$ for some $\omega_{ij} \in \{0, 1\}$,

$$\langle \overline{e_i}, \overline{f_i}, \overline{K_i}, \overline{K_i'} \rangle \simeq \mathcal{H}_{q,\omega}^+(\mathfrak{g})$$

Proof of Lemma

- Decomposition of \mathbf{f}_i, K'_i follows from explicit calculation using Feigin's embedding.
- Decomposition of \mathbf{e}_i, K_i requires combinatorics of Coxeter moves:

Lemma (I. (2020))

If $l(s_i w s_j) = l(w)$, then there is a sequence of Coxeter moves that brings the reduced word of $w \in W$:

$$\mathbf{i} = (i, \dots) \mapsto \mathbf{i}' = (\dots, j)$$

where the sequence of Coxeter moves splits into 2 stages, the second of which increase in indices consecutively from first to last letter.

Proof of Lemma

- Decomposition of \mathbf{f}_i, K'_i follows from explicit calculation using Feigin's embedding.
- Decomposition of \mathbf{e}_i, K_i requires combinatorics of Coxeter moves:

Lemma (I. (2020))

If $l(s_i w s_j) = l(w)$, then there is a sequence of Coxeter moves that brings the reduced word of $w \in W$:

$$\mathbf{i} = (i, \dots) \mapsto \mathbf{i}' = (\dots, j)$$

where the sequence of Coxeter moves splits into 2 stages, the second of which increase in indices consecutively from first to last letter.

Proof of Lemma

- Decomposition of \mathbf{f}_i, K'_i follows from explicit calculation using Feigin's embedding.
- Decomposition of \mathbf{e}_i, K_i requires combinatorics of Coxeter moves:

Lemma (I. (2020))

If $l(s_i w s_j) = l(w)$, then there is a sequence of Coxeter moves that brings the reduced word of $w \in W$:

$$\mathbf{i} = (i, \dots) \mapsto \mathbf{i}' = (\dots, j)$$

where the sequence of Coxeter moves splits into 2 stages, the second of which increase in indices consecutively from first to last letter.

Proof of Lemma

Example

$$\mathfrak{g} = \mathfrak{sl}_5: \mathbf{i} = (1, 2, 1, 3, 4, 3, 2, 3, 1, 2) \rightsquigarrow (\dots\dots\dots, 4)?$$

Stage 1:

$$(1, 2, 1, 3, 4, 3, 2, 3, 1, 2) \rightsquigarrow (1, 2, 1, 3, 2, 1, 4, 3, 2, 1)$$

Stage 2:

$$\begin{aligned} & (1, 2, 1, 3, 2, 1, 4, 3, 2, 1) \\ & \rightsquigarrow (2, 1, 2, 3, 2, 1, 4, 3, 2, 1) \\ & \rightsquigarrow (2, 1, 3, 2, 3, 1, 4, 3, 2, 1) \\ & \rightsquigarrow (2, 1, 3, 2, 1, 3, 4, 3, 2, 1) \\ & \rightsquigarrow (2, 1, 3, 2, 1, 4, 3, 4, 2, 1) \\ & \rightsquigarrow (2, 1, 3, 2, 1, 4, 3, 2, 4, 1) \\ & \rightsquigarrow (2, 1, 3, 2, 1, 4, 3, 2, 1, 4) \end{aligned}$$

Proof of Lemma

Example


$\mathfrak{g} = \mathfrak{sl}_5$: $\mathbf{i} = (1, 2, 1, 3, 4, 3, 2, 3, 1, 2) \rightsquigarrow (\dots\dots\dots, 4)$?

Stage 1:


$$(1, 2, 1, 3, 4, 3, 2, 3, 1, 2) \rightsquigarrow (1, 2, 1, 3, 2, 1, 4, 3, 2, 1)$$

Stage 2:


$$\begin{aligned} & (1, \underline{2, 1}, 3, 2, 1, 4, 3, 2, 1) \\ & \rightsquigarrow (2, 1, \underline{2, 3, 2}, 1, 4, 3, 2, 1) \\ & \rightsquigarrow (2, 1, 3, 2, \underline{3, 1}, 4, 3, 2, 1) \\ & \rightsquigarrow (2, 1, 3, 2, 1, \underline{3, 4, 3}, 2, 1) \\ & \rightsquigarrow (2, 1, 3, 2, 1, 4, 3, \underline{4, 2}, 1) \\ & \rightsquigarrow (2, 1, 3, 2, 1, 4, 3, 2, \underline{4, 1}) \\ & \rightsquigarrow (2, 1, 3, 2, 1, 4, 3, 2, 1, 4) \end{aligned}$$

Example: E_6 

$$A_1 \subset A_2 \subset A_3 \subset D_4 \subset D_5 \subset E_6$$

Example: E_6 

$$D_5 \subset E_6$$

Example: B_4 

$$J = \{1, 2, 3\} \subset I = \{1, 2, 3, 4\}, \quad 1 = \text{short}$$

Further Discussions

- Lusztig's braid group action T_i as cluster mutations on $\mathbf{Q}(\bar{\mathbf{i}})$

$$\mathbf{e}_i \longleftrightarrow \mathbf{f}_i^*, \quad \mathbf{f}_i \longleftrightarrow \mathbf{e}_i^*$$

- Geometric meaning of the cluster structure of $\mathbf{D}(\bar{\mathbf{i}})$
 - partial configuration space $\text{Conf}_w^e(\mathcal{A})$. [Goncharov-Shen]
- Combinatorial description of $\mathcal{U}_q(\mathfrak{g}) \hookrightarrow \mathbf{X}_q^{\mathbf{Q}}$
 - $\pi(\mathbf{e}_i), \pi(\mathbf{f}_i)$ are polynomials in X_i , not Laurent.
 - Type A_n : counting of cycles in dual plabic graphs.
- Tensor product decompositions of $\mathcal{P}_\lambda^J \otimes \mathcal{P}_{\lambda'}^{J'}$
 - \mathcal{R} matrix well-defined \implies new braided tensor category?
 - Study the spectrum of Casimir operators \mathbf{C}_k .
 - Proved for $J = \emptyset$ and $\mathfrak{g} = \mathfrak{sl}_{n+1}$. [Schrader-Shapiro]
- Generalization to other modules of affine quantum groups $\mathcal{U}_q(\widehat{\mathfrak{g}}_{\mathbb{R}})$?

Further Discussions

- Lusztig's braid group action T_i as cluster mutations on $\mathbf{Q}(\bar{\mathbf{i}})$

$$\mathbf{e}_i \longleftrightarrow \mathbf{f}_i^*, \quad \mathbf{f}_i \longleftrightarrow \mathbf{e}_i^*$$

- Geometric meaning of the cluster structure of $\mathbf{D}(\bar{\mathbf{i}})$
 - partial configuration space $\text{Conf}_w^e(\mathcal{A})$. [Goncharov-Shen]
- Combinatorial description of $\mathcal{U}_q(\mathfrak{g}) \hookrightarrow \mathbf{X}_q^{\mathbf{Q}}?$
 - $\pi(\mathbf{e}_i), \pi(\mathbf{f}_i)$ are polynomials in X_i , not Laurent.
 - Type A_n : counting of cycles in dual plabic graphs.
- Tensor product decompositions of $\mathcal{P}_\lambda^J \otimes \mathcal{P}_{\lambda'}^{J'}$
 - \mathcal{R} matrix well-defined \implies new braided tensor category?
 - Study the spectrum of Casimir operators \mathbf{C}_k .
 - Proved for $J = \emptyset$ and $\mathfrak{g} = \mathfrak{sl}_{n+1}$. [Schrader-Shapiro]
- Generalization to other modules of affine quantum groups $\mathcal{U}_q(\widehat{\mathfrak{g}}_{\mathbb{R}})?$

Further Discussions

- Lusztig's braid group action T_i as cluster mutations on $\mathbf{Q}(\bar{\mathbf{i}})$

$$\mathbf{e}_i \longleftrightarrow \mathbf{f}_{i^*}, \quad \mathbf{f}_i \longleftrightarrow \mathbf{e}_{i^*}$$

- Geometric meaning of the cluster structure of $\mathbf{D}(\bar{\mathbf{i}})$
 - partial configuration space $\text{Conf}_w^e(\mathcal{A})$. [Goncharov-Shen]
- Combinatorial description of $\mathcal{U}_q(\mathfrak{g}) \hookrightarrow \mathbf{X}_q^{\mathbf{Q}}$
 - $\pi(\mathbf{e}_i), \pi(\mathbf{f}_i)$ are polynomials in X_i , not Laurent.
 - Type A_n : counting of cycles in dual plabic graphs.
- Tensor product decompositions of $\mathcal{P}_\lambda^J \otimes \mathcal{P}_{\lambda'}^{J'}$
 - \mathcal{R} matrix well-defined \implies new braided tensor category?
 - Study the spectrum of Casimir operators \mathbf{C}_k .
 - Proved for $J = \emptyset$ and $\mathfrak{g} = \mathfrak{sl}_{n+1}$. [Schrader-Shapiro]
- Generalization to other modules of affine quantum groups $\mathcal{U}_q(\widehat{\mathfrak{g}}_{\mathbb{R}})$?

Further Discussions

- Lusztig's braid group action T_i as cluster mutations on $\mathbf{Q}(\bar{\mathbf{i}})$

$$\mathbf{e}_i \longleftrightarrow \mathbf{f}_i^*, \quad \mathbf{f}_i \longleftrightarrow \mathbf{e}_i^*$$

- Geometric meaning of the cluster structure of $\mathbf{D}(\bar{\mathbf{i}})$
 - partial configuration space $\text{Conf}_w^e(\mathcal{A})$. [Goncharov-Shen]
- Combinatorial description of $\mathcal{U}_q(\mathfrak{g}) \hookrightarrow \mathbf{X}_q^{\mathbf{Q}}$
 - $\pi(\mathbf{e}_i), \pi(\mathbf{f}_i)$ are polynomials in X_i , not Laurent.
 - Type A_n : counting of cycles in dual plabic graphs.
- Tensor product decompositions of $\mathcal{P}_\lambda^J \otimes \mathcal{P}_{\lambda'}^{J'}$
 - \mathcal{R} matrix well-defined \implies new braided tensor category?
 - Study the spectrum of Casimir operators \mathbf{C}_k .
 - Proved for $J = \emptyset$ and $\mathfrak{g} = \mathfrak{sl}_{n+1}$. [Schrader-Shapiro]
- Generalization to other modules of affine quantum groups $\mathcal{U}_q(\widehat{\mathfrak{g}}_{\mathbb{R}})$?

Further Discussions


- Lusztig's braid group action T_i as cluster mutations on $\mathbf{Q}(\bar{\mathbf{i}})$

$$\mathbf{e}_i \longleftrightarrow \mathbf{f}_i^*, \quad \mathbf{f}_i \longleftrightarrow \mathbf{e}_i^*$$

- Geometric meaning of the cluster structure of $\mathbf{D}(\bar{\mathbf{i}})$
 - partial configuration space $\text{Conf}_w^e(\mathcal{A})$. [Goncharov-Shen]
- Combinatorial description of $\mathcal{U}_q(\mathfrak{g}) \hookrightarrow \mathbf{X}_q^{\mathbf{Q}}$
 - $\pi(\mathbf{e}_i), \pi(\mathbf{f}_i)$ are polynomials in X_i , not Laurent.
 - Type A_n : counting of cycles in dual plabic graphs.
- Tensor product decompositions of $\mathcal{P}_\lambda^J \otimes \mathcal{P}_{\lambda'}^{J'}$
 - \mathcal{R} matrix well-defined \implies new braided tensor category?
 - Study the spectrum of Casimir operators \mathbf{C}_k .
 - Proved for $J = \emptyset$ and $\mathfrak{g} = \mathfrak{sl}_{n+1}$. [Schrader-Shapiro]
- Generalization to other modules of affine quantum groups $\mathcal{U}_q(\widehat{\mathfrak{g}}_{\mathbb{R}})$?


Thank you for your attention!

Half-Dehn Twist



4 flips

=



$$R = \mu_1 \cdots \mu_n$$

acting as unitary
operator on $P_\lambda \otimes P_\mu$.
 $\cong L^2(\mathbb{R}^{2N})$